



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

or 6; but $s=6$ makes t impossible, while $s=5$ makes $t=4$, and we have $2st=40$ and $(s^2-t^2)=18$. So $2pq=80$ and $(p^2-q^2)=18$, as before. Take $m=50$, p may be 6 or 7, but 6 makes q impossible, while $p=7$ gives $q=1$, and $2pq=14$ and $p-q^2=48$. $\therefore 14^2+48^2=50^2$. But $50=5 \times 10$, and $5=4+1$; then $s=2$ and $t=1$, $2st=4$, and $s^2-t^2=3$, $2pq=40$, and $p^2-q^2=30$, and $30^2+40^2=50^2$. Solutions may be obtained from other factors, but they give one of these two results. If $m=65$, it may be shown in the same manner that there are 4 pairs of numbers answering the conditions.

II. Solution by R. J. ADCOCK, Larchland, Illinois.

Since $(x^2+y^2+z^2+u^2+v^2)^2=(x^2+y^2+z^2+u^2+v^2)^2+(2xv)^2+(2yv)^2+(2zv)^2+(2uv)^2$, is true for two or any greater number of letters. Then

$$\left[(2x+1)^2+(2y)^2 \right]^2 = \left[(2x+1)^2-(2y)^2 \right]^2 + \left[2(2x+1)2y \right]^2.$$

$$\text{And, } 1 = \left[\frac{(2x+1)^2-(2y)^2}{(2x+1)^2+(2y)^2} \right]^2 + \left[\frac{2(2x+1)2y}{(2x+1)^2+(2y)^2} \right]^2.$$

If n^2 be the given square number,

$$n^2 = \left[\frac{(2x+1)^2-(2y)^2}{(2x+1)^2+(2y)^2} \times n \right]^2 + \left[\frac{2(2x+1)2yn}{(2x+1)^2+(2y)^2} \right]^2 \text{ are the}$$

two square parts into which n^2 is divided. Observing that x may have any value including 0, y any value not = 0, and in order to avoid repetition of parts into which the square number is divided, the numbers for x and y must not make $2x+1$ and $2y$ have a common factor or be equal. If $x=0$, and $y=1$, the two square parts of n^2 are $\left(\frac{3n}{5}\right)^2$ and $\left(\frac{4n}{5}\right)^2$.

Also solved by H. W. Draughton, G. B. M. Zerr, P. S. Berg, A. L. Foote, P. H. Philbrick, and H. C. Whitaker.

2. Proposed by J. M. COLAW, Principal of High School, Monterey, Virginia.

Find two numbers, such that the difference of their squares may be a cube, and the difference of their cubes a square.

I. Solution by G. B. M. ZERR, A. M., Principal of High School, Staunton, Virginia.

Let $6x^3$, $10x^3$ be the numbers.

Then $100x^6-36x^6=64x^6$ = a cube number, and $1000x^9-216x^9=784x^9=(28)^2x^9$ = a square number, when x is a perfect square. Let $x=1$, then 6 and 10 are the numbers. Other values can be obtained by substituting 4, 9, 16, etc. for x .

II Solution by C. W. M. BLACK, Department of Mathematics in Wilmington Conference Academy, Dover, Delaware.

(A). Represent the numbers by x , and $x+a$. The difference of the cubes will be (1), $a(3x^2+3ax+a^2)$, and the difference of the squares (2), $a(2x+a)$. For one solution the first will be a square if $a=3x^2+3ax+a^2$, and the second a cube if $a=(a+2x)^2$. Solving these equations, $a=1$ and $x=-1$ or 0, $a=0$ and $x=0$, making the required numbers -1 and 0, 0 and 1, or 0 and 0. Also, $a(2x+a)$ will be a cube if (3), $a^2=a+2x$. Combining (1) and (3), $(a-1)(3a^2+3a+4)=0$. The first factor gives $a=1$, whence $x=-1$ or 0, as before. The second factor gives an imaginary result. As these results are not satisfactory, we seek another method for finding other values, if there are any.

(B). It may easily be shown that if x and $x+a$ are positive integers

and prime to each other, there is in general no solution. So, we have only to find how two numbers not prime to each other may be obtained to fulfill the requirements

(C). Let n and p be two positive integers prime to each other, then (4), $n^3 - p^3 = hr^2$, and (5), $n^2 - p^2 = ks^3$, where r, s, h, k , are positive integers. Let m be a factor such that if n and p are both multiplied by it, they will fulfill the required condition. Then, (6), $m^3 n^3 - m^3 p^3 = m^3 hr^2 =$ a square. (7), $m^2 n^2 - m^2 p^2 = m^2 ks^3 =$ a cube. The first will be a square if $m = hc^2$ (8), and the second a cube if $m = kc^3$ (9). Combining (8) and (9), $c = \frac{h}{k}$ and $m = \frac{h^3}{k^2}$ (10).

Then by taking any two positive integers prime to each other, substituting, and finding values of h and k in equations (4) and (5), and multiplying each member by the value of m from (10), we have two numbers which fulfill the requirements. There is nothing in above proof to limit the method to positive integers prime to each other, but these may be taken as the basis of all positive numbers. When one of the numbers taken is negative, different results are derived. When both are negative, the value derived for m will change the numbers so that the difference of the cubes will not be negative. (Ex. 1) $5^3 - 3^3 = 98 = 2 \times 7^2$, $5^2 - 3^2 = 16 = 2 \times 2^3$, $h=2, k=2, m=2$. \therefore numbers are 10 and 6. (Ex. 2). $2^3 - 1^3 = 7$, $2^2 - 1^2 = 3$, $h=7, k=3, m = \frac{7^3}{3^2}$. \therefore numbers are $\frac{686}{9}$ and $\frac{343}{9}$. (Ex. 3).

$2^3 - (-1)^3 = 9$, $2^2 - (-1)^2 = 3$, $h=1, k=3, m = \frac{1}{9}$. \therefore numbers are $\frac{2}{9}$ and $-\frac{1}{9}$.

(Ex. 4). $1^3 - 0^3 = 1$, $1^2 - 0^2 = 1$, $h=1, k=1, m=1$. \therefore numbers are 1 and 0, as derived before in (A).

Also solved by *H. W. Draughon, A. L. Foote, P. H. Philbrick, and G. B. M. Zerr.*

3. Proposed by O. S. KIBLER, Superintendent of Schools, West Middleburg, Logan County, Ohio.

It is required to find three whole numbers in an arithmetical progression, such that the sum of every two of them shall be a square.

Solution by A. L. FOOTE, 80 Broad St., New York City.

Let $\frac{1}{2}x^2 - y$, $\frac{1}{2}x^2$, and $\frac{1}{2}x^2 + y$ be the numbers. Then we must have x^2 , $x^2 + y$, and $x^2 - y$ each squares and since x^2 is a square, we require only to make $x^2 + y$ and $x^2 - y$ squares. Let $x^2 + y = m^2$ and $x^2 - y = n^2$. Let $y = 2sx + s^2$ and we have $x^2 + y = x^2 + 2sx + s^2 = (x+s)^2$, and so we have but to find $x^2 - 2sx - s^2 = n^2$. Now let $x^2 - 2sx - s^2 = (x-m)^2 = x^2 - 2mx + m^2$ and we have $x = \frac{m^2 + s^2}{2m - 2s}$

where m and s may be assumed at pleasure. Let $m=5$ and $s=4$, then

$x = \frac{25+16}{10-8} = \frac{41}{2}$ and $\frac{1}{2}x^2 = \frac{1681}{8}$, $y = 2sx + s^2 = 164 + 16 = 180$, so we have $\frac{1681}{8}$, $\frac{241}{8}$, and $\frac{3121}{8}$. To render these integral multiply each by 16 and we have

482, 3362 and 6242 for the required numbers. The squares are $3844 = 62^2$, $6724 = 82^2$, and $9604 = 98^2$. The values of m must be so chosen that $\frac{1}{2}x^2 - y$ will be positive.

J. H. Drummond finds 380, 8450 and 16514. Also solved by P. S. Berg and H. W. Draughon.